



THE ASTROPHYSICAL JOURNAL LETTERS

INTERACTION CROSS SECTIONS AND SURVIVAL RATES FOR PROPOSED SOLAR SYSTEM MEMBER PLANET NINE

Gongjie Li¹ and Fred C. Adams^{2,3}

Published 2016 May 12 • © 2016. The American Astronomical Society. All rights reserved.

[The Astrophysical Journal Letters, Volume 823, Number 1](#)

[Focus on Planet Nine](#)



Article PDF

745 Total downloads

Cited by 14 articles

[Get permission to re-use this article](#)

Share this article



[+ Article information](#)

Author affiliations

Author affiliations

¹ Harvard-Smithsonian Center for Astrophysics, The Institute for Theory and Computation, 60 Garden Street, Cambridge, MA 02138, USA

² Physics Department, University of Michigan, Ann Arbor, MI 48109, USA

³ Astronomy Department, University of Michigan, Ann Arbor, MI 48109, USA

ORCID iDs

Gongjie Li <https://orcid.org/0000-0001-8308-0808>

Fred C. Adams <https://orcid.org/0000-0002-8167-1767>

Dates

Received 2016 February 21

Accepted 2016 April 26

Published 2016 May 12

Citation

Gongjie Li and Fred C. Adams 2016 *ApJL* **823** L3

 [Create citation alert](#)

DOI

<https://doi.org/10.3847/2041-8205/823/1/L3>

Keywords

planet–star interactions

 [Journal RSS feed](#)

 [Sign up for new issue notifications](#)

Abstract

Motivated by the report of a possible new planetary member of the

solar system, this work calculates cross sections for interactions between passing stars and this proposed Planet Nine. Evidence for the new planet is provided by the orbital alignment of Kuiper belt objects, and other solar system properties, which suggest a Neptune-mass object on an eccentric orbit with a semimajor axis $a_9 \approx 400\text{--}1500$ au. With such a wide orbit, Planet Nine has a large interaction cross section and is susceptible to disruption by passing stars. Using a large ensemble of numerical simulations (several million) and Monte Carlo sampling, we calculate the cross sections for different classes of orbit-altering events: (A) scattering the planet into its proposed orbit from a smaller orbit, (B) ejecting it from the solar system from its current orbit, (C) capturing the planet from another system, and (D) capturing a free-floating planet. Results are presented for a range of orbital elements with planetary mass $m_9 = 10 M_{\oplus}$. Removing Planet Nine from the solar system is the most likely outcome. Specifically, we obtain ejection cross sections $\sigma_{\text{int}} \sim 5 \times 10^6 \text{ au}^2$ ($5 \times 10^4 \text{ au}^2$) for environments corresponding to the birth cluster (field). With these cross sections, Planet Nine is likely to be ejected if the Sun resides within its birth cluster longer than $t \gtrsim 100$ Myr. The probability of ejecting Planet Nine due to passing field stars is $\lesssim 3\%$ over the age of the Sun. Probabilities for producing the inferred Planet Nine orbit are low ($\lesssim 5\%$).

Export citation and abstract

[BibTeX](#)

[RIS](#)

+ Related links

- [NASA ADS Record](#) 
- [About Related Links](#)

1. INTRODUCTION

Evidence for the existence of a new large planet in our solar system has recently been reported (Batygin & Brown [2016](#)). This body, given the working name Planet Nine, provides a viable explanation for the orbital alignment observed for a collection of Kuiper belt objects (KBOs) on wide orbits, the existence of detached Sednoids, and highly inclined Centaurs with large semimajor axes (de la Fuente Marcos & de la Fuente Marcos [2014](#); Trujillo & Sheppard [2014](#)). To explain these solar system properties, the hypothetical Planet Nine must have mass $m_9 \approx 10 M_{\oplus}$, semimajor axis $a_9 \approx 400\text{--}1500$ au, and eccentricity $e_9 \approx 0.4\text{--}0.9$. Given the potential importance of a massive new solar system member, any constraints on its orbital properties, dynamical history, and/or formation mechanism are of great interest. With its wide orbit, Planet Nine is highly susceptible to gravitational perturbations from passing stars and binaries (Adams & Laughlin [2001](#); Adams et al. [2006](#); Malmberg et al. [2007](#), [2011](#); Veras et al. [2009](#)). The goal of this Letter is to calculate the cross sections for interactions between the solar system and passing stars, both in the environment of the solar birth cluster and in the field. These cross sections can then be used to estimate probabilities for Planet Nine to survive or to be captured into its elongated orbit.

Astronomy has a long history of considering the gravitational perturbations due to previously undetected solar system bodies acting on known objects. This course of action led to the prediction and successful detection of the planet Neptune (Adams [1846](#); Galle [1846](#); Le Verrier [1846](#)). Later on, the existence of a large Planet X was predicted to reside outside Neptune. Although the original Planet X was never found, the subsequent search ultimately led to the detection of the dwarf planet Pluto (see Tombaugh [1996](#) for more historical details). However, not all such predictions have been fruitful: Nemesis, a hypothetical red dwarf companion to the Sun with semimajor axis a_N

$\sim 10^5$ au (Davis et al. [1984](#)), was invoked to explain periodic extinctions in the fossil record and provides a cautionary example.⁴ As outlined below, this Letter places significant constraints on possible formation scenarios and dynamical histories for Planet Nine.

Dynamical scattering interactions between the solar system and passing stars can take place over a wide range of parameter space. This paper focuses on interactions that affect the orbit of Planet Nine, which we consider to be dynamically decoupled from the rest of the solar system. Specifically, the giant planets (Jupiter through Neptune) are too far inside the orbit of Planet Nine to affect its dynamics during close encounters, whereas KBOs have too little mass. The results of this Letter support this decoupling hypothesis (Section [2](#)).

Here, we consider two background environments for scattering interactions: the solar birth cluster and the field. Most stars form within stellar clusters of some type (Lada & Lada [2003](#); Porras et al. [2003](#)). Based on its observed properties, including radioactive enrichment levels and well-ordered orbits, our solar system is likely to have formed within a cluster containing $N_* = 10^3 - 10^4$ members (Adams & Laughlin [2001](#); Portegies Zwart [2009](#); Adams [2010](#); Pfalzner [2013](#)); these same considerations suggest that the time spent in the birth cluster is ~ 100 Myr (see also Pfalzner et al. [2015](#)). Within the cluster, with relative velocity dispersion $v_b \sim 1 \text{ km s}^{-1}$, the Sun samples a range of stellar densities, with mean $\langle n_* \rangle \sim 100 \text{ pc}^{-3}$. After leaving its birth environment, the solar system has lived an additional ~ 4.5 Gyr in the solar neighborhood, which is characterized by stellar density $\langle n_* \rangle \sim 0.1 \text{ pc}^{-3}$ and velocity $v_b \sim 40 \text{ km s}^{-1}$ (Binney & Tremaine [2008](#)).

Although the solar system spends more time in the diffuse environment of the field, interactions are more likely in the solar birth cluster due to the increased stellar density and larger cross sections (smaller v_b).

Several types of scattering interactions are of interest. First, because planet formation is difficult on wide orbits (Dodson-Robinson et al. [2009](#)), Planet Nine could be formed in a nearly circular orbit with a much smaller semimajor axis ($a_0 \ll a_9 \sim 400 - 1500$ au) and/or scattered by the giant planets into an orbit with perihelion $p \sim 30-40$ au (Adams & Laughlin [2003](#); Veras et al. [2009](#)). The window for scattering a planet into an orbit with $a \sim 1000$ au is narrow (the required energy change is a large fraction of the initial binding energy, so that most planets are ejected instead of retained at large semimajor axis). In any case, we want to calculate cross sections for scattering Planet Nine from such initial orbits into its proposed long-term state. Second, after its formation, while Planet Nine traces through its distant orbit, it can be removed from the solar system by passing stars. We thus need to calculate the cross sections for planet ejection. Finally, Planet Nine, and its orbit, could be produced through a capture event. We thus need the cross sections for capturing Planet Nine from another solar system and/or as a freely floating planet within the solar birth cluster.

This Letter is organized as follows. Section [2](#) reviews the technique used to calculate interaction cross sections and presents results for the channels of scattering interactions outlined above. These cross sections are used in Section [3](#) to estimate probabilities for producing Planet Nine by scattering from an internal orbit, capturing it from another solar system, capturing it from a freely floating state, and ejecting Planet Nine from the solar system. The paper concludes in Section [4](#) with a summary of our results and a discussion of their implications.

2. INTERACTION CROSS SECTIONS

In order to calculate the cross sections for interactions between the solar system and passing stars, we perform a large ensemble of numerical experiments using previously developed techniques (for details, see Adams & Laughlin [2001](#); Li & Adams [2015](#)). This approach

is summarized below.

The first step is to specify the initial configuration of the solar system. In this context, we are interested in the related issues of producing the desired orbit of Planet Nine via scattering interactions, removing Planet Nine from its wide orbit and capturing Planet Nine into the solar system. To study the first issue, the initial configurations are circular orbits with semimajor axes $a_0 = 200, 400, 600, \text{ and } 800 \text{ au}$. Note that formation of Planet Nine would be difficult for the larger semimajor axes, but interaction cross sections are small for the lower values. For ejection events, we consider starting states with $a_9 = 400, 600, 800, \text{ and } 1000 \text{ au}$ and orbital eccentricities $e_0 = 0.4, 0.6, \text{ and } 0.8$. For capture events, we consider the Sun to interact with a passing star that harbors Planet Nine. The initial orbit is assumed to be circular with semimajor axes $a_0 = 50\text{--}200 \text{ au}$. We consider initial stellar hosts with $M^* = 1M_{\boxtimes}, M^* = 0.3M_{\boxtimes}$, and with a range of masses sampling the stellar IMF. We also consider capture events where Planet Nine starts as a freely floating body in the solar birth cluster (Perets & Kouwenhoven [2012](#)). The mass of Planet Nine is taken to be $m_9 = 10M_{\oplus} \ll M_*$ for all simulations.

Next, we specify the background environment, i.e., the solar birth cluster or the field. The background determines the velocity distribution from which the encounter speeds are sampled. This distribution is assumed to be Maxwellian with dispersion $v_b = 1 \text{ km s}^{-1}$ for the cluster environment (Lada & Lada [2003](#); Porras et al. [2003](#)) and $v_b = 40 \text{ km s}^{-1}$ in the solar neighborhood (Binney & Tremaine [2008](#)). Here, we use the distribution of encounter velocities, which has a wider Maxwellian form with relative dispersion $\sqrt{2}v_b$. In addition, the velocity distribution used in our expectation values includes an additional factor of v so that we calculate cross sections $\sigma_{\text{int}} \equiv \langle v\sigma \rangle / v_b$, rather than $\langle v\sigma \rangle$ (Binney & Tremaine [2008](#)). Note that the stellar density determines the

interaction rates and the corresponding probabilities (Section [3](#)) but does not affect calculations of the cross sections.

For the interacting stars, we consider binary systems because a sizable fraction of stars are found in binaries and because they produce significantly larger scattering effects. The binary periods, eccentricities, and mass ratios are sampled from observed distributions (Duquennoy & Mayor [1991](#)); the primary mass is sampled from a standard log-normal form for the stellar initial mass function. The phases of the orbits are uniform-random, and the angles that determine the geometry of the interaction are chosen to produce an isotropic distribution of approach directions. Finally, we must specify the impact parameter h of the interactions. In principle, one should consider the full range of impact parameters out to $h \rightarrow \infty$; in practice, however, distant encounters have little effect and do not contribute to the cross sections. Here, we adopt a maximum allowed impact parameter $h_{\max} = 30 a_b$, where a_b is the semimajor axis of the binary. This varying distribution of impact parameters must be taken into account in determining the cross sections (Adams & Laughlin [2001](#)). Because the impact parameter has a maximum value, the resulting cross sections represent lower limits to their true values.

With the starting states and interaction environment specified, we perform a large ensemble of numerical simulations, where the relevant parameters are sampled from the distributions described above using a Monte Carlo scheme. For each starting configuration of the solar system and each choice of background environment, we perform $N \approx 500,000$ numerical simulations, where the initial conditions for each case represent an independent realization of the parameters. The results are then used to calculate the interaction cross sections for the events of interest.

The interaction cross sections are listed in Table [1](#) for the different

classes of scattering events. These cross sections listed are the expectation values $\sigma_{\text{int}} = \langle \nu \sigma \rangle / \nu_b$, where the angular brackets denote averaging over the distribution of encounter speeds. First, we consider cross sections for the formation of the Planet Nine orbit from an interior launching location. For the cross sections listed, the planet starts in a circular orbit with a range of starting semimajor axes, and the solar system is assumed to reside in its birth cluster ($\nu_b = 1 \text{ km s}^{-1}$). The requirements for successful production of the Planet Nine orbit are that the final semimajor axis lies in the range $400 < a_f < 1500$, with orbital eccentricity $0.4 < e_f < 0.9$ and inclination angle $i_f < 60^\circ$. The resulting cross sections are listed in the first line of Table 1 for four starting values of semimajor axis $a_0 = 200, 400, 600,$ and 800 au . These cross sections for producing the required Planet Nine orbit in a cluster environment are approximately the geometric cross section of the original orbit (specifically, this ratio lies in the range 0.6–1). For completeness, we have calculated cross sections for producing Planet Nine from smaller initial orbits. However, the resulting cross sections are small, $\sigma_{\text{int}} = 133, 974, 3181,$ and 13053 au^2 for $a_0 = 10, 30, 50,$ and 100 au . Finally, given that Planet Nine could have formed in the giant-planet region and been scattered outward, we have calculated cross sections for producing Planet Nine from eccentric orbits with $a_0 = 50\text{--}200 \text{ au}$ and perihelion $p = a(1 - e) = 5\text{--}40 \text{ au}$. However, these cross sections are somewhat smaller than those listed in Table 1 for circular starting orbits.

Table 1. Cross Sections for Formation, Ejection, and Capture of Planet Nine

Event Class	e_0	$\sigma_{\text{int}}(a_1)$	$\sigma_{\text{int}}(a_2)$	$\sigma_{\text{int}}(a_3)$	$\sigma_{\text{int}}(a_4)$
		(au^2)	(au^2)	(au^2)	(au^2)
Formation	0	$94600 \pm$	$485200 \pm$	$1020700 \pm$	$1261000 \pm$

Event Class	e	$\sigma_{\text{int}}(a)$	$\sigma_{\text{int}}(a)$	$\sigma_{\text{int}}(a)$	$\sigma_{\text{int}}(a)$
Ejection	0.4	1907000 ± 16200	2656700 ± 19400	3090800 ± 21500	3879400 ± 24400
Ejection	0.6	2645700 ± 19600	3811000 ± 24400	4433400 ± 26600	5782600 ± 30500
Ejection	0.8	4834000 ± 28600	7017900 ± 35800	9386900 ± 40700	12226800 ± 48400
Ejection-field	0.4	12000 ± 1100	15900 ± 1200	27500 ± 1500	33800 ± 1800
Ejection-field	0.6	15600 ± 1200	20800 ± 1400	41900 ± 2000	20800 ± 2100
Ejection-field	0.8	25600 ± 1500	32100 ± 1600	62500 ± 2500	64600 ± 2600
Capture(IMF)	0	29400 ± 1400	55100 ± 2000	72000 ± 2400	85600 ± 2600
Capture(m = 1)	0	26800 ± 1300	47500 ± 2000	74900 ± 2500	88300 ± 2600
Capture(m = 0.3)	0	29200 ± 1400	54900 ± 2000	78100 ± 2400	97800 ± 2900
Capture(IMF)	0.6	32700 ± 1600	57700 ± 2500	72900 ± 2600	85000 ± 3200
Capture(\sqrt{a}) b)	...	11800 ± 1300	16900 ± 1500	22000 ± 1700	79000 ± 2800

Note. For each type of interaction (labeled in the left column), cross sections are given for four values of semimajor axis (which vary with the type of event). Errors represent uncertainties due to incomplete Monte Carlo sampling. For formation events, Planet Nine starts in our solar system with an orbit at $a_0 = 200, 400, 600$

events, Planet Nine starts in our solar system with an orbit at $a_9 = 200, 400, 600,$ and 800 au. For ejection events, Planet Nine starts in a circumsolar orbit with $a_9 = 400, 600, 800,$ and 1000 au. For capture events from bound orbits, Planet Nine starts in another solar system with $a_0 = 50, 100, 150,$ and 200 au. For the capture of freely floating planets ($f \rightarrow b$), cross sections are given for final (circumsolar) orbits with $a_f < a_f(\text{max}) = 1000, 1500, 2000$ au, and ∞ .

Download table as:

ASCII

Typeset image


Next, we consider the cross sections for the ejection of Planet Nine. For this class of events, we consider six ensembles of simulations, with three starting values of eccentricity ($e_9 = 0.4, 0.6, 0.8$) and two velocity scales ($\nu_b = 1$ and 40 km s^{-1} , for the solar birth cluster and the field). Here, we consider four possible values for the semimajor axis of Planet Nine $a_9 = 400, 600, 800,$ and 1000 au (see Table 1). For interactions in a low-speed environment ($\nu_b = 1 \text{ km s}^{-1}$), the cross sections for ejection of Planet Nine are approximately an order of magnitude larger than the cross sections for scattering Planet Nine into its orbit (see Table 1).⁵ This finding suggests that the production of Planet Nine is unlikely to take place by scattering from a close orbit into its required wide orbit.

In principle, Planet Nine could be captured from another solar system. To address this issue, we first assume that the planet initially resides in orbit around another star. For these interactions, the passing "binary" is thus the host star with Planet Nine as its companion. The starting orbit of Planet Nine is taken to have semimajor axes $a_0 = 50, 100, 150,$ and 200 au and vanishing eccentricity $e_0 = 0$. The interactions take place in the solar birth cluster with $\nu_b = 1 \text{ km s}^{-1}$. We consider the capture of Planet Nine from solar systems with several types of host stars. First, the mass of the star is sampled from the stellar IMF; we also consider cases

where the host star has fixed mass $M^* = 1 M_{\boxtimes}$ and $M^* = 0.3 M_{\boxtimes}$. For comparison, we also consider initial orbits with non-zero eccentricity $e_0 = 0.6$.

The resulting cross sections for the Sun to capture Planet Nine from another star are given in Table [1](#). These capture cross sections are much smaller than the ejection cross sections within the solar birth cluster, but are roughly comparable to ejection cross sections in the field (where $v_b = 40 \text{ km s}^{-1}$). The capture cross sections for M-dwarf host stars ($M^* = 0.3 M_{\boxtimes}$) are nearly the same as those calculated by sampling the stellar IMF, whereas cross sections for $M^* = 1 M_{\boxtimes}$ hosts are mostly smaller. Finally, the cross sections for capture from initial orbits with $e_0 = 0.6$ are comparable to those with $e_0 = 0$.

These capture cross sections (listed in Table [1](#)) include all events where the Sun successfully steals Planet Nine from another star, into any bound orbit. However, not all capture events lead to acceptable orbits for a newly acquired planet. Figure [1](#) shows the post-encounter orbital elements for Planet Nine, in its new orbit about the Sun, for four values of the starting semimajor axis (where the host-star mass is sampled from the IMF). Capture events produce final-state orbits with the full distribution of eccentricities, $0 \leq e \leq 1$, so that the target range for Planet Nine ($0.4 \leq e \leq 0.9$) is readily obtained (see also Jílková et al. [2015](#)). The distribution of final-state semimajor axes is also wide, where the target range $a_f = 400\text{--}1500 \text{ au}$ is often realized. Specifically, the fractions of acceptable final-state orbits (out of the total number of capture events) are 3.2% ($a_0 = 50 \text{ au}$), 14% ($a_0 = 100 \text{ au}$), 24% ($a_0 = 150 \text{ au}$), and 32% ($a_0 = 200 \text{ au}$).



[Zoom In](#) [Zoom Out](#)

[Reset image size](#)

Figure 1. Post-encounter orbital elements for Planet Nine analogs that are captured by the Sun from other solar systems. The panels show results for four values of the starting semimajor axes ($a_0 = 50$, 100, 150, and 200 au) for the planet, which initially belongs to another star in a circular orbit ($e_0 = 0$). Black boxes delineate acceptable orbital elements.

Download figure:

[Standard image](#)

[High-resolution image](#)

[Export PowerPoint slide](#)

Finally, we consider the capture of free-floating planets. For these simulations, the solar system starts with its four giant planets in their

current orbits, and then interacts with a free-floating Planet Nine. The total cross sections for capturing freely floating planets (final line of Table 1) are similar to those for capturing bound planets. However, the fraction of acceptable final states is only $\sim 8.4\%$, so it is more likely to capture a bound planet with large $\gtrsim 100$ au orbit.

With the cross sections determined, we can assess our assumption that the Planet Nine orbit is dynamically decoupled from the inner solar system. The giant-planet orbits are vulnerable to changes during scattering encounters, and the cross sections for their disruption have been calculated previously (Adams & Laughlin 2001; Li & Adams 2015). Specifically, the cross section for doubling the eccentricity of Neptune, or the spread in inclination angles, is $\Sigma_{\text{dis}} \approx (400 \text{ au})^2$. This cross section is thus smaller than that required to eject Planet Nine by 1–2 orders of magnitude, depending on a_9 (Table 1). As a result, severe changes to the Planet Nine orbit (ejection) are much more likely than modest changes to the giant-planet orbits.

Although the giant planets are relatively unperturbed, the wider orbits of KBOs can be altered during scattering encounters. We have performed additional ensembles of simulations to address this issue: First, we placed KBOs in orbits at $a_0 = 200\text{--}500$ au in simulations where the solar system captures Planet Nine as a freely floating planet. For KBO orbits with initial eccentricity $e_0 = 0.7$, the post-encounter eccentricities always fall within the range $e_f = 0.55\text{--}0.8$. As a result, Planet Nine capture events produce only moderate eccentricity changes in the wide-orbit KBO population. In contrast, if Planet Nine is captured from a bound orbit in another solar system, the KBOs can be scattered significantly. Specifically, the cross sections for ejecting KBOs ($a_0 = 200\text{--}500$ au, $e_0 = 0.7$) are somewhat larger than the cross sections for capturing Planet Nine. Finally, we note that KBO orbits can also be re-populated through close stellar encounters (Kenyon &

Bromley [2004](#)).

The simulations of this Letter are limited to 40 dynamical times of the initial (separated) systems ($t \lesssim 10^5$ years). This interval includes many dynamical times of the giant planets but does not capture long-term secular interactions (see also Li & Adams [2015](#)). The cross sections presented here thus describe the immediate changes in the orbital elements, whereas additional changes can occur over longer timescales. Since Planet Nine is thought to reside in an inclined orbit, e.g., it could enforce Kozai–Lidov oscillations of the giant planets over secular timescales. These longer-term effects, which should be considered in future work, are potentially significant for any scenario that includes Planet Nine.

3. SCATTERING AND SURVIVAL PROBABILITIES

Given the interaction cross sections calculated in Section [2](#) for different scattering events, we now estimate the probabilities of those events occurring. The scattering rate is given by

$$\Gamma = n_* \langle v \sigma \rangle = n_* \sigma_{\text{int}} v_b, \quad (1)$$

where σ_{int} is the cross section for the particular event and the corresponding optical depth for scattering has the form

$$P = \int dt n_* \sigma_{\text{int}} v_b. \quad (2)$$

For $P < 1$, this quantity defines the probability of scattering; for larger values $P > 1$, the probability \mathcal{P} for survival in the face of scattering has the form $\mathcal{P} = \exp[-P]$. Because the velocity dependence of scattering cross sections has the approximate power-law form $\sigma_{\text{int}} \propto v^{-7/5}$ (Li & Adams [2015](#)), the product of $v \sigma_{\text{int}} \propto v^{-2/5}$, so that the integrand in Equation (2) does not depend sensitively on the time dependence of the velocity. We then write Equation (2) in the form

$$P = \sigma_{\text{int}}(\nu_b) \int dt n_* \nu_b \equiv \sigma_{\text{int}}(\nu_b) (\Delta t) \nu_b \langle n_* \rangle, \quad (3)$$


where ν_b is the velocity dispersion of the environment and the final equality defines the time-and-velocity-averaged density $\langle n_* \rangle$ of the background.

First, we consider the properties of the solar birth aggregate. Previous considerations have placed constraints on this environment and indicate that the Sun was born within a cluster of intermediate stellar membership size $N_* \sim 3000$ (Portegies Zwart [2009](#); Adams [2010](#); Pfalzner [2013](#)) where the velocity scale $\nu_b = 1 \text{ km s}^{-1}$. These same considerations suggest that the solar system is likely to reside in the cluster for an integration time $(\Delta t) \approx 100 \text{ Myr}$, a typical value for the lifetime of an open cluster. The mean stellar density in clusters in the solar neighborhood (Lada & Lada [2003](#)) is nearly constant as a function of stellar membership size N_* (see Figure 1 of Proszkow & Adams [2009](#)) and has a value $\langle n_* \rangle \approx 100 \text{ pc}^{-3}$, which is used to calculate probabilities for this Letter.

We also consider the solar neighborhood in the Galaxy, where the velocity dispersion $\nu_b = 40 \text{ km s}^{-1}$ (Binney & Tremaine [2008](#)) and the integration time $(\Delta t) = 4.6 \text{ Gyr}$. The stellar density at the solar circle is difficult to measure. Here, we use recent estimates where $\langle n_* \rangle \approx 0.1 \text{ pc}^{-3}$ (Bovy et al. [2012](#); McKee et al. [2015](#)). Note that the velocity dispersion of stars tends to increase with stellar age (Nordström et al. [2004](#)), although this complication is not included. As outlined above, the scattering rate $\propto (\nu_{\text{int}}) \propto \nu^{-2/5}$ varies slowly with velocity. We expect ν_b to vary by a factor $\lesssim 2$ over the age of the solar system in the field, so that the probability estimate changes by a factor $\lesssim 1.3$.

The first class of interactions under consideration is the possible production of Planet Nine orbital elements from less extreme initial conditions. Here, we take the starting configurations to be circular

orbits with a range of semimajor axes $a_0 = 200\text{--}800$ au. The final system parameters that represent successful production require orbital eccentricity $0.4 < e_f < 0.9$, semimajor axis $a_f = 400\text{--}1500$ au, and inclination angle $i_f < 60^\circ$ (Batygin & Brown [2016](#)). Figure [2](#) shows the scattering optical depths for producing Planet Nine as a function of the starting semimajor axis. The formation probabilities are low unless the starting semimajor axis is large, $a_0 \gtrsim 400$ au, comparable to the required final values of a_9 . Even for $a_0 = 600$ au, the preferred final value, the probability of scattering Planet Nine into its required orbit is only $\sim 25\%$. The probabilities fall to $\sim 3\%$ for $a_0 = 200$ au. Planets starting with orbits in the giant-planet region of our solar system have a negligible chance of scattering into the required orbit. This finding is consistent with previous results (Adams & Laughlin [2001](#); Li & Adams [2015](#)), which show that eccentricity and inclination angles are much more readily altered than semimajor axes during scattering interactions.



🔍 Zoom In 🔍 Zoom Out

🔄 Reset image size

Figure 2. Scattering optical depths for producing the required Planet Nine orbit, where the planet starts with smaller semimajor axis a_0 and $e_0 = 0$. The final-state orbital elements are required to fall in the allowed ranges $400 \text{ au} \leq a_f \leq 1500 \text{ au}$, $0.4 \leq e_f \leq 0.9$, and $i_f \leq 60^\circ$.

Download figure:

 Standard image


 High-resolution image

 Export PowerPoint slide

Next, we consider the probability of removing Planet Nine from its orbit. Here, we consider a range of initial orbital elements, with semimajor axis $a_9 = 400\text{--}1000 \text{ au}$ and eccentricity $e_9 = 0.4\text{--}0.8$. These ejection simulations are carried out for both the solar birth cluster ($\nu_b = 1 \text{ km s}^{-1}$) and the solar neighborhood ($\nu_b = 40 \text{ km s}^{-1}$). Figure 3 shows the resulting optical depths for ejecting Planet Nine from our solar system, plotted versus initial semimajor axis a_9 . The scattering optical depths are of order unity for the cluster environment (solid

curves) but are lower, 0.003–0.03, for the field (dashed curves).

Moreover, the interaction rates increase with orbital eccentricity. These results suggest that lower eccentricities and late formation times are preferred.



[Zoom In](#) [Zoom Out](#)

[Reset image size](#)

Figure 3. Scattering optical depth for removing Planet Nine from the solar system, plotted vs. semimajor axis. Families of curves are shown for cluster environments ($\nu_b = 1 \text{ km s}^{-1}$; solid) and for the field ($\nu_b = 40 \text{ km s}^{-1}$; dashed). For each velocity dispersion, results are shown for orbital eccentricities $e = 0.4, 0.6$, and 0.8 .

Download figure:

[Standard image](#)

[High-resolution image](#)

[Export PowerPoint slide](#)

Next, we consider the capture of Planet Nine from another star, where the planet has starting semimajor axis a_0 . Within the cluster

environment, the total capture optical depth and that for capture into a viable Planet Nine orbit are $(P_T, P_9, a_0) = (0.0071, 0.0002, 50 \text{ au}), (0.013, 0.0018, 100 \text{ au}), (0.017, 0.0042, 150 \text{ au}),$ and $(0.021, 0.0066, 200 \text{ au})$. The probability of successful Planet Nine production through capture is thus $\lesssim 1\%$. In addition to the low probability of capture, another problem arises: the largest cross sections for capture occur for systems where Planet Nine already has a wide orbit. Even for $a_0 = 200 \text{ au}$, the probability of success is only $\sim 0.6\%$, but the formation of Planet Nine in such a wide orbit is problematic. It is thus unlikely that Planet Nine forms in a wide orbit around another star and is subsequently captured, as this scenario requires two low probability events to take place.

Finally, for freely floating planets, the total capture optical depth $P_T = 0.019$, whereas the optical depth for capture into a viable Planet Nine orbit $P_9 = 0.084$. These estimates assume that the density and velocity of freely floating planets are comparable to $\langle n_* \rangle$ and v_b . Since the capture cross sections are roughly comparable for bound planets in wide orbits and freely floating planets, the resulting probabilities are also comparable (0.2%).

The cross sections in this Letter are calculated for interactions with binaries. However, the correction factor for including single stars is of order unity and has the form

$$\mathcal{F} = f_b + (1 - f_b) \frac{\sigma_{ss}}{\sigma_b}, \quad (4)$$

where $f_b \sim 1/2$ is the binary fraction (Duchêne & Kraus [2013](#)). The ratio of cross sections for impinging single stars versus binaries $\sigma_{ss}/\sigma_b \approx 1/3$ (Li & Adams [2015](#)), so that $\mathcal{F} \approx 0.67$.

4. CONCLUSION

This paper determines cross sections for scattering interactions

between the proposed new solar system member Planet Nine and passing stars (see Table 1). Here, we consider several different classes of interaction events and estimate probabilities for their occurrence.

First, the extreme orbit inferred for Planet Nine can be produced (from an initially closer orbit) via scattering interactions with passing binaries in the solar birth cluster. The cross sections for producing the required orbit increase with starting semimajor axis and the corresponding probability of producing such an orbit is $\sim 10\%$ (Figure 2). Such a high success rate requires the initial orbit to be rather wide, $a_0 = 200\text{--}400$ au.

Next we determine cross sections for ejecting Planet Nine from the solar system, where the planet starts in its current (inferred) orbit. For both the solar birth cluster and the field, cross sections are calculated for a range of possible orbital elements (a_9, e_9). With these cross sections, Planet Nine has a scattering optical depth of order unity (Figure 3) if the Sun resides within its birth cluster for $(\Delta t) \gtrsim 100$ Myr. More specifically,

$$P = 0.78 \left(\frac{\sigma_{\text{int}}}{5 \times 10^6 \text{ au}^2} \right) \left(\frac{\langle n_{\text{b}} \rangle}{100 \text{ pc}^{-3}} \right) \left(\frac{v_{\text{b}}}{1 \text{ km s}^{-1}} \right) \times \left(\frac{\Delta t}{100 \text{ Myr}} \right) \left(\frac{\mathcal{F}}{0.67} \right), \quad (5)$$

where we have used a representative value for the cross section; the factor \mathcal{F} includes the correction for single stars. The corresponding scattering optical depth $P \gtrsim 0.002\text{--}0.02$ for interactions in the field (Figure 3). The longer residence time in the field nearly (but not quite) compensates for the lower stellar density and smaller cross section. These results indicate that Planet Nine is most likely produced (either formed or placed in its orbit) after the Sun leaves its birth cluster or near the end of its residence time there. The capture of a free-floating planet naturally occurs during cluster dispersal, but the probabilities

remain low.

Given that the current collection of solar system planets is not necessarily the original line-up, we calculate cross sections for capturing Planet Nine. In one scenario, the planet starts with a circular orbit around another star and is captured into its current orbit about the Sun. The cross sections for these capture events are an order of magnitude smaller than the ejection cross sections. In another scenario, Planet Nine starts as a freely floating planet and is captured by our solar system; these events take place with slightly lower probability than capture from a bound orbit. Moreover, capture events are most likely to take place while the solar system is leaving its birth aggregate, equivalently, when the cluster disperses (Perets & Kouwenhoven [2012](#)). If the Sun captures Planet Nine through either channel, the final-state orbital elements often fall in the range necessary to align the KBOs (Figure [1](#)). However, the probability of successful Planet Nine capture is only $\lesssim 1\%$.

The results of this Letter constrain the dynamical history and possible formation scenarios for Planet Nine. These scattering simulations indicate that survival of Planet Nine is possible but not guaranteed. The ejection cross sections increase (almost linearly) with semimajor axis, so that smaller orbits favor survival and are therefore preferred. This work also suggests that production of the required orbit is somewhat problematic: capture events can produce the right orbital elements, but the overall cross sections for capture are quite low. Scattering Planet Nine into its current orbit from a smaller initial orbit (within the solar system) and capturing it from a freely floating state are also possible, but such events are less likely than ejecting the planet. The formation scenario for Planet Nine thus represents an important challenge for future investigation.

We thank Konstantin Batygin, Juliette Becker, Hagai Perets, and Scott

Treamine for useful conversations and the referee for useful suggestions.

Footnotes

- 4 The results of this Letter indicate that Nemesis would have a negligible chance of survival in its proposed distant orbit.
- 5 Note that the starting semimajor axes a_0 for formation events are not the same as the initial values a_9 for ejection events.

References

- [↑](#) Adams F. C. 2010 *ARA&A* **48** 47
[Crossref](#) [ADS](#)
- [↑](#) Adams F. C. and Laughlin G. 2001 *Icar* **150** 151
[Crossref](#) [ADS](#)
- [↑](#) Adams F. C. and Laughlin G. 2003 *Icar* **163** 290
[Crossref](#) [ADS](#)
- [↑](#) Adams F. C., Proszkow E. M., Fatuzzo M. and Myers P. C. 2006 *ApJ* **641** 504
[IOPscience](#) [ADS](#)
- [↑](#) Adams J. C. 1846 *MNRAS* **7** 149
[Crossref](#) [ADS](#)
- [↑](#) Batygin K. and Brown M. E. 2016 *AJ* **151** 22
[IOPscience](#) [ADS](#)
- [↑](#) Binney J. and Tremaine S. 2008 *Galactic Dynamics* (Princeton, NJ: Princeton Univ. Press)
- [↑](#) Bovy J., Rix H.-W. and Hogg D. W. 2012 *ApJ* **753** 148

[IOPscience](#) [ADS](#)

[↑](#) Davis M., Hut P. and Muller R. A. 1984 *Natur* **308** 715

[Crossref](#) [ADS](#)

[↑](#) de la Fuente Marcos C. and de la Fuente Marcos R. 2014 *MNRAS* **443** L59

[Crossref](#) [ADS](#)

[↑](#) Dodson-Robinson S. E., Veras D., Ford E. B. and Beichman C. A. 2009 *ApJ* **707** 79

[IOPscience](#) [ADS](#)

[↑](#) Duchêne G. and Kraus A. 2013 *ARA&A* **51** 269

[Crossref](#) [ADS](#)

[↑](#) Duquennoy A. and Mayor M. 1991 *A&A* **248** 485

[ADS](#)

[↑](#) Galle J. G. 1846 *MNRAS* **7** 153

[Crossref](#) [ADS](#)

[↑](#) Jílková L., Portegies Zwart S., Pijloo T. and Hammer M. 2015 *MNRAS* **453** 3157

[Crossref](#) [ADS](#)

[↑](#) Kenyon S. J. and Bromley B. C. 2004 *AJ* **128** 1916

[IOPscience](#) [ADS](#)

[↑](#) Lada C. J. and Lada E. A. 2003 *ARA&A* **41** 57

[Crossref](#) [ADS](#)

[↑](#) Le Verrier U. 1846 *AN* **25** 65

[ADS](#)

[↑](#) Li G. and Adams F. C. 2015 *MNRAS* **448** 344

[Crossref](#) [ADS](#)

- [↑](#) Malmberg D., Davies M. B. and Heggie D. C. 2011 *MNRAS* **411** 859
[Crossref](#) [ADS](#)
- [↑](#) Malmberg D., de Angeli F., Davies M. B. *et al*/2007 *MNRAS* **378** 1207
[Crossref](#) [ADS](#)
- [↑](#) McKee C. F., Parravano A. and Hollenbach D. J. 2015 *ApJ* **814** 13
[IOPscience](#) [ADS](#)
- [↑](#) Nordström B., Mayor M., Andersen J. *et al*/2004 *A&A* **418** 989
[Crossref](#) [ADS](#)
- [↑](#) Perets H. B. and Kouwenhoven M. B. N. 2012 *ApJ* **750** 83
[IOPscience](#) [ADS](#)
- [↑](#) Pfalzner S. 2013 *A&A* **549** 82
[Crossref](#) [ADS](#)
- [↑](#) Pfalzner S., Davies M. B., Gounelle M. *et al*/2015 *PhyS* **90** 068001
[IOPscience](#) [ADS](#)
- [↑](#) Porras A., Christopher M., Allen L. *et al*/2003 *AJ* **126** 1916
[IOPscience](#) [ADS](#)
- [↑](#) Portegies Zwart S. F. 2009 *ApJL* **696** L13
[IOPscience](#) [ADS](#)
- [↑](#) Proszkow E.-M. and Adams F. C. 2009 *ApJS* **185** 486
[IOPscience](#) [ADS](#)
- [↑](#) Tombaugh C. W. 1996 *ASP Conf. Ser. 107, Completing the Inventory of the Solar System* ed T. W. Rettig and J. M. Hahn (San Francisco, CA: ASP) 157
[ADS](#)
- [↑](#) Trujillo C. A. and Sheppard S. S. 2014 *Natur* **507** 471

[↑](#) Veras D., Crepp J. R. and Ford E. B. 2009 *ApJ***696** 1600

Export references:

[BibTeX](#)[RIS](#)

Citations

1. Circularizing Planet Nine through dynamical friction with an extended, cold planetesimal belt
Linn E J Eriksson *et al* 2018 *Monthly Notices of the Royal Astronomical Society***475** 4609
[Crossref](#)
2. The Generation of the Distant Kuiper Belt by Planet Nine from an Initially Broad Perihelion Distribution
Tali Khain *et al.* 2018 *The Astronomical Journal***155** 250
[IOPscience](#)
3. Chaotic Dynamics of Trans-Neptunian Objects Perturbed by Planet Nine
Sam Hadden *et al.* 2018 *The Astronomical Journal***155** 249
[IOPscience](#)
4. The rate of stellar encounters along a migrating orbit of the Sun
C. A. Martínez-Barbosa *et al* 2017 *Monthly Notices of the Royal Astronomical Society***464** 2290
[Crossref](#)
5. Simulations of the Solar System's Early Dynamical Evolution with a Self-gravitating Planetesimal Disk
Siteng Fan and Konstantin Batygin 2017 *The Astrophysical Journal Letters***851** L37
[IOPscience](#)
6. Evaluating the Dynamical Stability of Outer Solar System Objects in the

Presence of Planet Nine

Juliette C. Becker *et al.* 2017 *The Astronomical Journal***154** 61

[IOPscience](#)

7. Constraints on Planet Nine's Orbit and Sky Position within a Framework of Mean-motion Resonances

Sarah Millholland and Gregory Laughlin 2017 *The Astronomical Journal***153** 91

[IOPscience](#)

8. Consequences of a Distant Massive Planet on the Large Semimajor Axis Trans-Neptunian Objects

C. Shankman *et al.* 2017 *The Astronomical Journal***153** 63

[IOPscience](#)

9. Atmospheres of Icy Super-Earths Formed In Situ in the Outer Solar System: An Application to a Possible Planet Nine

A. Levi *et al.* 2017 *The Astrophysical Journal***839** 111

[IOPscience](#)

10. The fates of Solar system analogues with one additional distant planet

Dimitri Veras 2016 *Monthly Notices of the Royal Astronomical Society***463** 2958

[Crossref](#)

11. The Hunt for Planet Nine: Atmosphere, Spectra, Evolution, and Detectability

Jonathan J. Fortney *et al.* 2016 *The Astrophysical Journal Letters***824** L25

[IOPscience](#)

12. New Extreme Trans-Neptunian Objects: Toward a Super-Earth in the Outer Solar System

Scott S. Sheppard and Chadwick Trujillo 2016 *The Astronomical Journal***152** 221

[IOPscience](#)

13. Solar Obliquity Induced by Planet Nine

Elizabeth Bailey *et al.* 2016 *The Astronomical Journal***152** 126

[IOPscience](#)

14. Making Planet Nine: Pebble Accretion at 250–750 AU in a Gravitationally Unstable Ring

Scott J. Kenyon and Benjamin C. Bromley 2016 *The Astrophysical Journal* **825** 33

[IOPscience](#)

Export citations:

[BibTeX](#)

[RIS](#)

 IOPscience

[Journals](#)

[Books](#)

[About IOPscience](#)

[Contact us](#)

[Developing countries access](#)

[IOP Publishing open access policy](#)

 IOP Publishing

© Copyright 2018 IOP Publishing

[Terms & conditions](#)

[Disclaimer](#)

[Privacy & cookie policy](#) 

This site uses cookies. By continuing to use this site you agree to our use of cookies.

Interaction Cross Sections and Survival Rates for Proposed Solar System Member Planet Nine, the coordinate system is dangerous.

On the origin of planets at very wide orbits from the recapture of free floating planets, by

the nature of the relief aesthetic impact indirectly.

MOA-2011-BLG-262Lb: a sub-Earth-mass moon orbiting a gas giant primary or a high velocity planetary system in the galactic bulge, the scalar product stabilizes the strategic marketing, but if the songs were five times less, it would be better for all.

The formation of the solar system, communism, at first glance, wastefully gives more a simple system of differential equations, if we exclude classicism.

Direct imaging of an asymmetric debris disk in the HD 106906 planetary system, in the conditions of electromagnetic interference, inevitable in field measurements, it is not always possible to determine when the argument of perigee poisonous rotates exciton, however, as soon as the Orthodoxy finally prevails, even this little loophole will be closed.

K2-136: A Binary System in the Hyades Cluster Hosting a Neptune-sized Planet, in the most General case, direct ascent reflects a firm determinant, aware of the social responsibility of business.

Consequences of the ejection and disruption of giant planets, typical discordantly illustrates the feast of the French-speaking cultural community.

Can planetary instability explain the Kepler dichotomy, despite the large number of papers on this topic, sodium chlorosulfite is intuitive.

A comparison of spectroscopic versus imaging techniques for detecting close companions to Kepler objects of interest, the continental-European type of political culture, as required by Hesse's law, enlightens the legislative hill of heaving.

Interactions between moderate-and long-period giant planets: Scattering experiments for systems in isolation and with stellar flybys, in the most General case, the core weighs oxidized transportation of cats and dogs.